§520.530

§ 520.530 Cythioate oral liquid.

- (a) *Specifications*. Each milliliter contains 15 milligrams of cythioate.
- (b) *Sponsor*. See Nos. 000859 and 053501 in §510.600 of this chapter.
- (c) Special considerations. Cythioate is a cholinesterase inhibitor. Do not use this product in animals simultaneously with or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, insecticides, pesticides, or chemicals.
- (d) Conditions of use—(1) Amount. 15 milligrams cythicate per 10 pounds of body weight every third day or twice a week.
- (2) Indications for use. Dogs, for control of fleas.
- (3) *Limitations.* For oral use in dogs only. Do not use in greyhounds or in animals that are pregnant, sick, under stress, or recovering from surgery. Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- [49 FR 5614, Feb. 14, 1984, as amended at 67 FR 78355, Dec. 24, 2002]

§520.531 Cythioate tablets.

- (a) [Reserved]
- (b) *Sponsors.* See No. 000859 in §510.600(c) of this chapter for use of 30-and 90-milligram (mg) tablets and see No. 053501 in §510.600(c) of this chapter for use of 30-mg tablet.
- (c) Special considerations. Cythioate is a cholinesterase inhibitor. Do not use this product in animals simultaneously with or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, insecticides, pesticides, or chemicals.
- (d) Conditions of use—(1) Amount. 30 milligrams cythicate per 20 pounds of body weight every third day or twice a week.
- (2) Indications for use. Dogs, for control of fleas.
- (3) *Limitations.* For oral use in dogs only. Do not use in greyhounds or in animals that are pregnant, sick, under stress, or recovering from surgery. Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- [49 FR 5615, Feb. 14, 1984, as amended at 59 FR 26942, May 25, 1994; 67 FR 78355, Dec. 24, 2002]

§520.534 Decoquinate.

- (a) *Specifications*. The drug is a powder containing 0.8 percent decoquinate.
- (b) Sponsor. See No. 046573 in $\S 510.600(c)$ of this chapter.
- (c) Related tolerances. See §556.170 of this chapter.
- (d) Conditions of use. Calves—(1) Amount. Feed 22.7 milligrams per 100 pounds of body weight (0.5 milligram per kilogram) per day.
- (2) Indications for use. For the prevention of coccidiosis in ruminating and nonruminating calves, including veal calves, caused by *Eimeria bovis* and *E. zuernii*.
- (3) *Limitations*. Feed in whole milk at the rate of 22.7 milligrams per 100 pounds body weight daily (0.5 milligram per kilogram) for at least 28 days.
- [64 FR 10103, Mar. 2, 1999, as amended at 64 FR 30386, June 8, 1999]

§520.538 Deracoxib.

- (a) *Specifications*. Each chewable tablet contains 25 or 100 milligrams (mg) deracoxib.
- (b) Sponsor. See No. 058198 in $\S510.600$ (c) of this chapter.
 - (c) [Reserved]
- (d) Conditions of use in dogs—(1) Amount. Administer orally as needed, as a single daily dose based on body weight.
- (i) 1 to 2 mg/kilograms (kg) (0.45 to 0.91 mg/pound (lb), for use as in paragraph (d)(2)(i) of this section.
- (ii) 3 to 4 mg/kg (1.4 to 1.8 mg/lb) for up to 7 days, for use as in paragraph (d)(2)(ii) of this section.
- (2) *Indications for use.* (i) For the control of pain and inflammation associated with osteoarthritis.
- (ii) For the control of postoperative pain and inflammation associated with orthopedic surgery in dogs weighing 4 or more pounds (1.8 kg).
- (3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- [67 FR 68760, Nov. 13, 2002, as amended at 68 FR 18882, Apr. 17, 2003]